

WARNING
TOXIC, ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

Foxtr[®]  **t**

Contains:

69 g/litre fenoxaprop-p-ethyl in the form of an oil in water emulsion.

Also contains 34.5 g/litre cloquintocet-mexyl as a safener and 390 g/litre hydrocarbon liquid

**For selective grass weed control in autumn and spring sown
Barley, Wheat, Perennial Ryegrass and Pasture.**

Registered pursuant to the ACVM Act 1997, No. P008030
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, Approval Code HSR100112
See www.epa.govt.nz for approval conditions.

 **Distributed by Zelam Ltd**
Hudson Rd, New Plymouth, NZ Tel: 06 755 9234

 **CHEMINOVA**
HELPING YOU GROW

PLEASE READ THE LABEL COMPLETELY BEFORE USE
 This product must be under control of an approved handler during use.
 A record must be kept of the use and disposal of this product.



HSNO Classifications: 6.1E, 6.3B, 6.5B, 6.9B, 9.1C, 9.2D

WARNING: May be harmful if swallowed and enters airways. Causes mild skin irritation. May cause an allergic skin reaction. May cause damage to organs through prolonged or repeated exposure. Harmful to aquatic life with long lasting effects. Harmful to the soil environment. Avoid release to the environment.

PRECAUTIONS: DO NOT breathe vapours.

PERSONAL PROTECTION

DO NOT eat, drink or smoke while using. Wear protective clothing such as waterproof hat, coat and trousers when using. Wear protective gloves. Remove protective clothing and wash hands and face thoroughly before meals and after work. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid skin contact or inhalation. Avoid contact with the eyes. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse.

FIRST AID: If medical advice is needed, have product container or label at hand. If swallowed immediately call the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. DO NOT induce vomiting. If skin contact occurs, get medical advice/attention. If on skin wash with plenty of soap and water. If skin irritation or rash occurs get medical advice/attention. Get medical advice/attention if you feel unwell.

STORAGE: Store in original container tightly closed and in a locked dry, cool, well ventilated area, away from foodstuffs. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of this product if it is older than this.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section. Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

Container Disposal: Triple rinse container and add residue to mixture. Burn if circumstances, especially wind direction, permit. Otherwise bury in a suitable land fill.

SHELF LIFE: When stored appropriately, this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

SAFETY DATA SHEET: If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at www.zelam.com.

WEEDS CONTROLLED: Wild Oats, Lesser Canary Grass (*Phalaris minor*), Yellow Bristle Grass.

RESISTANT WEEDS: Gnawed Canary Grass (*Phalaris paradoxa*), Annual and Hybrid Ryegrasses, Tall Fescue, Chewing's Fescue, Cocksfoot, Browntop, Grazing Brome (=Spiky Brome – *Bromus stamineus*), Prairie Grass and most broadleaved weeds are not controlled by Foxtrot at label rates.

DIRECTIONS FOR USE:

CROP	WEEDS	RATE	REMARKS
Barley, Wheat	Wild Oats	750 mL/hectare in 250 litres of water. Add adjuvant oil at recommended rates	Apply when wild oats are between 3 leaf and 1st node visible growth stage and when they are growing actively
	Lesser Canary Grass (<i>Phalaris minor</i>)		Apply when weeds are between the 3 leaf visible stage and the end of tillering and are growing actively. Avoid application during periods of moisture stress.
Perennial ryegrass (<i>Lolium perenne</i>)	Wild Oats	750 mL/hectare in 250 litres of water. DO NOT use with any other agrochemicals or adjuvants.	Apply in early spring when wild oats are between 3 leaf and end of tillering growth stage and growing actively. Note: Seed yield response is likely to be reduced the longer the delay in the removal of wild oats in spring.
Pasture	Yellow Bristle Grass	Use 750 ml/ha in 250 litres of water. DO NOT use with any other agrichemicals or adjuvants.	Apply to established pasture 7 days after grazing during the period December/January.

MIXING: Shake well before use. Add the required amount to a partly filled spray tank with agitation system operating. Complete filling the tank with water and apply.

COMPATIBILITY: DO NOT USE FOXTROT IN ANY MIXTURES WHEN APPLYING TO PERENNIAL RYEGRASS. When used on barley and wheat, Foxtrot can be tankmixed with Cycocel* and Opus* in 2 and 3 way mixes. Care must be taken to ensure that recommended timings for all products are followed. The addition of a suitable adjuvant oil such as D-C-Trate or Hasten at label rates is recommended. DO NOT use Foxtrot within 5 days of broad-leaved herbicide applications.

APPLICATION: DO NOT overlap swath width in ryegrass crops as yield suppression may occur. Flush spray system with clean water after use.

RAINFASTNESS: Foxtrot is rainfast within one hour of application

WITHHOLDING PERIODS:

IT IS AN OFFENCE for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

Barley and Wheat: 70 days (DO NOT graze treated crop for at least 70 days after treatment)

Perennial Ryegrass: 42 days (DO NOT graze treated crop for at least 42 days after treatment)

Pasture: 28 days (Do not graze treated pasture for at least 28 days after treatment)

SOIL ACTIVITY: Foxtrot is rapidly broken down by soil microbes causing no restrictions on crop rotations.

UNDERSOWN CROPS: Foxtrot can be used on crops undersown with certain grasses (See under "Resistant Weeds" above) and clover. DO NOT use on crops undersown with "Concord" ryegrass.

MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY: Cheminova (NZ) Limited has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Cheminova (NZ) Limited accepts no responsibility for any loss arising from incorrect storage, handling or use.

Registered to Cheminova (NZ) Limited,
 141 Powderham St, New Plymouth City Centre, New Plymouth, Taranaki 4310 New Zealand
 © Registered trademark of Cheminova

* Other trademarks



EMERGENCY PHONE:
 0800 CHEMCALL – 0800 243 6225

Batch No:
 Date of Manufacture:



130402-10B